arXiv:math/0212367v1 [math.FA] 29 Dec 2002

Disintegration of cylindrical measures

Jan Pachl
pachl@acm.org

December 28, 2002*

Abstract

We show that the existence of disintegration for cylindrical measures follows from a general disintegration theorem for countably additive measures.

1 Notation

A *probability space* $(X, \mathcal{A}, P)$ is a nonempty set X together with a sigma-algebra $\mathcal{A}$ on X and a probability P on $\mathcal{A}$ (that is, a nonnegative countably additive measure with $PX = 1$). When $\mathcal{C}$ is a set of subsets of a given set, $\sigma(\mathcal{C})$ is the smallest sigma-algebra that contains $\mathcal{C}$. When $\mathcal{A}$ and $\mathcal{B}$ are sigma-algebras on X and Y, respectively, $\sigma(\mathcal{A} \otimes \mathcal{B})$ is the smallest sigma-algebra making the canonical projections $\pi_X : X \times Y \to X$ and $\pi_Y : X \times Y \to Y$ measurable.

When $(X,\ \mathcal{A},\ P)$ and $(Y,\ \mathcal{B},\ Q)$ are two probability spaces, a probability S on $\sigma(\mathcal{A} \otimes \mathcal{B})$ is a *joint probability* if $\pi_X[S] = P$ and $\pi_Y[S] = Q$.

A *lattice on* X is a class of subsets of X that is closed under finite unions and finite intersections.

A class $\mathcal{K}$ of sets is *semicompact* if every countable class $\mathcal{K}_0 \subseteq \mathcal{K}$ such that $\bigcap \mathcal{K}_0 = \emptyset$ contains a finite class $\mathcal{K}_{00} \subseteq \mathcal{K}_0$ such that $\bigcap \mathcal{K}_{00} = \emptyset$.

---

[*] Extracted from the author's manuscript dated July 20, 1979.

Let $(X, \mathcal{A}, P)$ be a probability space and $\mathcal{K} \subseteq \mathcal{A}$. We say that $\mathcal{K}$ *approximates* P if for every $E \in \mathcal{A}$ and $\varepsilon > 0$ there is a $K \in \mathcal{K}$ such that $K \subseteq E$ and $P(E \setminus K) < \varepsilon$.

All linear spaces are assumed to be over the field $\mathbf{R}$ of reals. When Y and Z are locally convex spaces, $\mathcal{L}(Y, Z)$ is the set of continuous linear mappings from Y to Z.

When I and J are two index sets such that $J \subseteq I$, the canonical projection from $\mathbf{R}^I$ onto $\mathbf{R}^J$ is denoted p_{IJ}.

When I is an infinite index set, let $\mathcal{C}(I)$ be the set of all subsets of $\mathbf{R}^I$ of the form $p_{IJ}^{-1}(C)$ where J is finite, $\emptyset \neq J \subseteq I$ and C is a compact subset of $\mathbf{R}^J$. Then $\mathcal{C}(I)_\delta$ is the set of all countable intersections of sets in $\mathcal{C}(I)$.

2 Disintegration theorem

The following theorem is an immediate consequence of Theorem 3.5 in [4].

Theorem 1 *Let $(X, \mathcal{A}, P)$ and $(Y, \mathcal{B}, Q)$ be two probability spaces, and let S be a joint probability on $\sigma(\mathcal{A} \otimes \mathcal{B})$. Let Q be complete, and let $\mathcal{K}$ be a semicompact lattice closed under countable intersections and such that $\mathcal{A} = \sigma(\mathcal{K})$ and $\mathcal{K}$ approximates P.*

Then there exists a family of probabilities $\{P_y\}_{y \in Y}$ on $\mathcal{A}$ such that

(a) for every $E \in \mathcal{A}$, the function $y \mapsto P_y E$ is $\mathcal{B}$-measurable;

(b) for every $E \in \mathcal{A}$ and $F \in \mathcal{B}$ we have

$$S(E \times F) \;=\; \int_F P_y E \;\mathrm{d}Q(y).$$

3 Application to cylindrical measures

In his study of non-linear images of cylindrical measures, Krée [1] raised the problem of the disintegration of cylindrical measures. Here we show that the disintegration in the sense of *4.A.b* in [1] always exists. However, the disintegration constructed here need not be continuous or measurable in the sense of *4.A.c-e* in [1]. In fact, the remark on page 36 in [2] seems to imply that there is an example, due to Schwartz, of a cylindrical measure with no measurable disintegration.

The following theorem improves Proposition (34) in [1]. The terminology and notation are as in [1] and [5].

Theorem 2 *Let Y and Z be two locally convex spaces and let m be a cylindrical probability on $Y \times Z$ whose canonical image m_1 on Y is countably additive. Extend m_1 to the complete probability Q defined on a sigma-algebra $\mathcal{B}$.*

Then there exists a family $\{m_y\}_{y\in Y}$ of cylindrical probabilities on Z such that for every positive integer n, every $b \in \mathcal{L}(Z, \mathbf{R}^n)$ and every Borel set $E \subseteq \mathbf{R}^n$,

(a) the function $y \mapsto b[m_y]E$ is $\mathcal{B}$-measurable on Y;

(b) for every $F \in \mathcal{B}$ we have

$$(I_Y \times b)[m](F \times E) = \int_F b[m_y]E \, \mathrm{d}Q(y).$$

The proof of the theorem uses the following result about projective limits of measures. The original version of this result is due to Bochner. For a proof of a more general result, see e.g. [3].

Theorem 3 *Let I be an infinite index set. Then*
(a) $\mathcal{C}(I)_\delta$ is a semicompact class of subsets of $\mathbf{R}^I$;
(b) if m is a nonnegative finitely additive measure on the algebra generated by $\mathcal{C}(I)$ and the image $p_{IJ}[m]$ in R^J is countably additive for each finite $J \subseteq I$ then m is countably additive and its unique extension to a countably additive measure on the sigma-algebra $\sigma(\mathcal{C}(I))$ is approximated by $\mathcal{C}(I)_\delta$.

Proof of Theorem 2. Schwartz ([5], pp. 177-180) points out that cylindrical measures on Z are in one-to-one correspondence with cylindrical measures on $\mathbf{R}^I$ for some index set I. Namely, Z is embedded in the algebraic dual Z'^*

of its topological dual Z', and Z'^* is isomorphic to $\mathbf{R}^I$ when I is chosen of the same cardinality as an algebraic basis of Z'. This yields an injective continuous linear mapping $h : Z \to \mathbf{R}^I$ such that $\mu \mapsto h[\mu]$ is a one-to-one correspondence between cylindrical probabilities μ on Z and cylindrical probabilities on $\mathbf{R}^I$. In the following we choose one such h and keep it fixed.

By Theorem 3, every cylindrical probability on $\mathbf{R}^I$ is countably additive, and we have a one-to-one correspondence between cylindrical probabilities on Z and countably additive probabilities on the sigma-algebra $\sigma(\mathcal{C}(I))$ in $\mathbf{R}^I$.

Consider the cylindrical probability $m_0 = (I_Y \times h)[m]$ on the space $Y \times \mathbf{R}^I$ and its projections on Y and $\mathbf{R}^I$. The projection $\pi_Y[m_0] = m_1$ is countably additive by the assumption. The projection $\pi_{\mathbf{R}^I}$ is countably additive as shown above; let P be its unique countably additive extension to the sigma-algebra $\mathcal{A} = \sigma(\mathcal{C}(I))$.

By a result of Marczewski [3], m_0 is countably additive. Thus m_0 has a unique extension to a countably additive probability S on the sigma-algebra $\sigma(\mathcal{A} \otimes \mathcal{B})$.

Hence we can apply Theorem 1 with $X = \mathbf{R}^I$ and $\mathcal{A}$, P, Y, $\mathcal{B}$, Q and S as defined. For each probability P_y on $\mathcal{A}$ obtained from Theorem 1 let m_y be the unique cylindrical probability on Z such that $h[m_y] = P_y$.

Since R^I identifies with Z'^*, every $b \in \mathcal{L}(Z, \mathbf{R}^n)$ factors through h, and it is now straightforward to verify the properties of m_y stated in the theorem.

Remark. Note that the proof does not use the fact that Y is a vector space. Thus the same result holds true, with the same proof, in the more general case of Y-cylindrical probabilities ([2], p. 34), for a topological or measurable space Y.

References

[1] P. Krée. Équations linéaires á coefficients aléatoires. *Symposia Mathematica Vol. VII*, pp. 515-546. Academic Press 1971.

[2] P. Krée. Accouplement de processus linéaires. Images de probabilités cylindriques par certaines applications non-linéaires. *Séminaire Goulaouic-Schwartz 1971-72.* Ecole Polytechnique Paris.

[3] E. Marczewski. On compact measures. *Fund. Math. 40* (1953), 113-124.

[4] J. Pachl. Disintegration and compact measures. *Math. Scand. 43* (1978), 157-168.

[5] L. Schwartz. Radon measures on arbitrary topological spaces and cylindrical measures. Oxford University Press 1973.